arXiv:math/0503666v1 [math.AC] 29 Mar 2005

SPECIAL SIMPLICES AND GORENSTEIN TORIC RINGS

HIDEFUMI OHSUGI AND TAKAYUKI HIBI

Abstract. Christos Athanasiadis [2] studies an effective technique to show that Gorenstein sequences coming from compressed polytopes are unimodal. In the present paper we will use such the technique to find a rich class of Gorenstein toric rings with unimodal h-vectors arising from finite graphs.

Introduction

Let $\mathcal{P} \subset \mathbb{R}^N$ be an integral convex polytope, i.e., a convex polytope each of whose vertices has integer coordinates. Let $K[\mathbf{x}, \mathbf{x}^{-1}, t] = K[x_1, x_1^{-1}, \ldots, x_N, x_N^{-1}, t]$ denote the Laurent polynomial ring in $(N+1)$ variables over a field K. The *toric ring* of $\mathcal{P}$ is the subalgebra $K[\mathcal{P}]$ of $K[\mathbf{x}, \mathbf{x}^{-1}, t]$ generated by those Laurent polynomials $\mathbf{x}^{\mathbf{a}}t = x_1^{a_1} \cdots x_N^{a_N} t$ such that $\mathbf{a} = (a_1, \ldots, a_N)$ is a vertex of $\mathcal{P}$. We will regard $K[\mathcal{P}]$ as a homogeneous algebra [3, p. 147] by setting each $\deg \mathbf{x}^{\mathbf{a}}t = 1$ and write $F(K[\mathcal{P}], \lambda)$ for its Hilbert series. One has $F(K[\mathcal{P}], \lambda) = (h_0 + h_1\lambda + \cdots + h_s\lambda^s)/(1-\lambda)^{d+1}$, where each $h_i \in \mathbb{Z}$ with $h_s \neq 0$ and where d is the dimension of $\mathcal{P}$. The sequence $(h_0, h_1, \ldots, h_s)$ is said to be the *h-vector* of $K[\mathcal{P}]$. If the toric ring $K[\mathcal{P}]$ is normal, then $K[\mathcal{P}]$ is Cohen–Macaulay. If $K[\mathcal{P}]$ is Cohen–Macaulay, then the h-vector of $K[\mathcal{P}]$ is nonnegative, i.e., each $h_i \geq 0$. Moreover, if $K[\mathcal{P}]$ is Gorenstein, then the h-vector of $K[\mathcal{P}]$ is symmetric, i.e., $h_i = h_{s-i}$ for all i.

An outstanding conjecture (which is still open) is that the h-vector of a Gorenstein toric ring is unimodal, i.e., $h_0 \leq h_1 \leq \cdots \leq h_{[s/2]}$. One of the established techniques to show that the h-vector $(h_0, h_1, \ldots, h_s)$ of a Gorenstein toric ring $K[\mathcal{P}]$ is unimodal is to find a simplicial convex polytope [14] of dimension $s-1$ whose h-vector coincides with $(h_0, h_1, \ldots, h_s)$. On the other hand, however, given a Gorenstein toric ring $K[\mathcal{P}]$, it seems difficult to find such a simplicial convex polytope.

Christos Athanasiadis [2] introduces the concept of a special simplex of a convex polytope. Let $\mathcal{P} \subset \mathbb{R}^N$ be a convex polytope. A $(q-1)$-simplex Σ each of whose vertices is a vertex of $\mathcal{P}$ is said to be a *special simplex* in $\mathcal{P}$ if each facet of $\mathcal{P}$ contains exactly $q-1$ of the vertices of Σ. Recall that an integral convex polytope $\mathcal{P} \subset \mathbb{R}^N$ is *compressed* [15, p. 337] (and [11]) if all "pulling triangulations" of $\mathcal{P}$ are unimodular. The toric ring $K[\mathcal{P}]$ of a compressed polytope $\mathcal{P}$ is normal. It turns out [2, Theorem 3.5] that if $\mathcal{P}$ is compressed and if there is a special simplex in $\mathcal{P}$, then the h-vector of $K[\mathcal{P}]$ is equal to the h-vector of a simplicial convex polytope.

In the present paper we will use [2, Theorem 3.5] to study the h-vector of the toric ring of the edge polytope of a finite graph satisfying the odd cycle condition as well as that of the stable polytope of a perfect graph.

1. Two polytopes arising from finite graphs

Let G be a finite graph on the vertex set $[n] = \{1, 2, \ldots, n\}$ having no loops and no multiple edges, and $E(G)$ the edge set of G. We associate each subset $W \subset [n]$ with the $(0,1)$-vector $\rho(W) = \sum_{j \in W} \mathbf{e}_j \in \mathbb{R}^n$. Here $\mathbf{e}_j$ is the j-th unit coordinate vector in $\mathbb{R}^n$. Thus in particular $\rho(\emptyset)$ is the origin of $\mathbb{R}^n$. A subset $W \subset [n]$ is called *stable* (resp. a *clique*) if $\{i, j\} \notin E(G)$ (resp. $\{i, j\} \in E(G)$) for all $i, j \in W$ with $i \neq j$. Note that the empty set as well as each single-element susbset of $[n]$ is both stable and a clique. Let $S(G)$ denote the set of stable sets of G.

We now introduce two convex polytopes arising from a finite graph G on $[n]$. First, the *edge polytope* [8] of G is the $(0,1)$-polytope $\mathcal{P}_G \subset \mathbb{R}^n$ which is the convex hull of $\{\rho(e) : e \in E(G)\}$. Second, the *stable polytope* [4] of G is the $(0,1)$-polytope $\mathcal{Q}_G \subset \mathbb{R}^n$ which is the convex hull of $\{\rho(W) : W \in S(G)\}$.

Example 1.1. Let P be a finite poset on $[n]$ and $\mathrm{com}(P)$ its comparability graph. Thus $\mathrm{com}(P)$ is the finite graph on $[n]$ such that $\{i, j\}$ with $i \neq j$ is an edge of $\mathrm{com}(P)$ if and only if i and j are comparable in P. Then the stable polytope of $\mathrm{com}(P)$ coincides with the chain polytope [16] of P.

The problem when the toric ring $K[\mathcal{P}_G]$ is normal and the problem when the edge polytope $\mathcal{P}_G$ possesses a unimodular covering [8, p. 420] are studied in [8] (and [13]).

Theorem 1.2 ([8]). *Given a finite connected graph G, the following conditions are equivalent:*

- (i) *The toric ring $K[\mathcal{P}_G]$ is normal;*
- (ii) *The edge polytope $\mathcal{P}_G$ possesses a unimodular covering;*
- (iii) *G satisfies the odd cycle condition, i.e., if each of C and C' is an odd cycle (a cycle of odd length) of G and if C and C' possess no common vertex, then there exists an edge $\{i, j\}$ of G such that i is a vertex of C and j is a vertex of C'.*

Thus in particular the edge polytope of a finite connected bipartite graph possesses a unimodular covering and its toric ring is normal.

A *chromatic number* of a finite graph G on $[n]$ is the smallest integer $\ell > 0$ for which there is a map $\varphi : [n] \to [\ell]$ with the property that $\varphi(i) \neq \varphi(j)$ if $\{i, j\} \in E(G)$. A finite graph G is called *perfect* if, for all induced subgraphs H of G including G itself, the chromatic number of H is equal to the maximal cardinality of cliques contained in H. The comparability graph of a finite partially ordered set is perfect ([1]).

The facets of the edge polytope $\mathcal{P}_G$ of a finite connected graph G is completely determined ([8, Theorem 1.7]). On the other hand, the facets of the stable polytope $\mathcal{Q}_G$ is completely determined when G is a perfect graph ([4, Theorem 3.1]).

2. Gorenstein toric rings

When $K[\mathcal{P}_G]$ (resp. $K[\mathcal{Q}_G]$) is normal, it follows easily that $K[\mathcal{P}_G]$ (resp. $K[\mathcal{Q}_G]$) coincides with the *Ehrhart ring* [6, p. 97] of $\mathcal{P}_G$ (resp. $\mathcal{Q}_G$). On the other hand, since the equations of the facets of $\mathcal{P}_G$ (resp. $\mathcal{Q}_G$) are known, when $K[\mathcal{P}_G]$ (resp.

$K[\mathcal{Q}_G]$) is normal, by using the criterion [5, Corollary (1.2)] one can determines the finite graphs G for which the toric ring $K[\mathcal{P}_G]$ (resp. $K[\mathcal{Q}_G]$) is Gorenstein.

Let G be a finite connected graph on $[n]$. Given a subset $V \neq \emptyset$ of $[n]$, write G_V for the induced subgraph of G on V. We say that G is 2-*connected* if G together with $G_{[n]\setminus\{i\}}$ for all $i \in [n]$ is connected. If $i \in [n]$, then $N(G;i)$ stands for the set of vertices j with $\{i,j\} \in E(G)$. If $T \subset [n]$, then $N(G;T) = \bigcup_{i \in T} N(G;i)$. The *bipartite graph induced by* a stable set $T \neq \emptyset$ of G is the bipartite graph on the vertex set $T \bigcup N(G;T)$ consisting of those edges $\{i,j\}$ of G with $i \in T$ and $j \in N(G;T)$.

Recall that a *matching* of G is a set of edges $\{e_1, \ldots, e_m\}$ such that $e_i \bigcap e_j = \emptyset$ for all $i \neq j$. A matching $\{e_1, \ldots, e_m\}$ of G is called *perfect* if $\bigcup_{i=1}^m e_i = [n]$. In particular n is even and $m = n/2$ if G possesses a perfect matching $\{e_1, \ldots, e_m\}$. It follows that G possesses a perfect matching if and only if the monomial $x_1 x_2 \cdots x_n$ belongs to the toric ring $K[\mathcal{P}_G]$.

Theorem 2.1. (a) *Let G be a finite connected graph on $[n]$ satisfying the odd cycle condition and suppose that every connected component of $G_{[n]\setminus\{i\}}$ possesses at least one odd cycle for all $i \in [n]$. Then the toric ring $K[\mathcal{P}_G]$ of the edge polytope $\mathcal{P}_G$ of G is Gorenstein if and only if (i) G possesses a perfect matching, (ii) one has $|N(G;T)| = |T| + 1$ for each stable set T of G such that the bipartite graph induced by T is connected with $T \bigcup N(G;T) \neq [n]$ and that every connected component of $G_{[n]\setminus(T \bigcup N(G;T))}$ has at least one odd cycle and (iii) one has $|T| = n/2 - 1$ for each stable set T of G such that the bipartite graph induced by T is connected with $T \bigcup N(G;T) = [n]$.*

(a') *Let G be a bipartite graph on $[n] = V_1 \bigcup V_2$ and suppose that G is 2-connected. Then the toric ring $K[\mathcal{P}_G]$ of the edge polytope $\mathcal{P}_G$ of G is Gorenstein if and only if (i) G possesses a perfect matching and (ii) one has $|N(G;T)| = |T| + 1$ for every subset $T \subset V_1$ such that $G_{T \bigcup N(G;T)}$ is connected and that $G_{[n]\setminus(T \bigcup N(G;T))}$ is a connected graph with at least one edge.*

(b) *The toric ring $K[\mathcal{Q}_G]$ of a stable polytope $\mathcal{Q}_G$ of a perfect graph G is Gorenstein if and only if all maximal cliques have the same cardinality.*

Proof. (a) The edge polytope $\mathcal{P}_G \subset \mathbb{R}^n$ lies on the hyperplane $\mathcal{H}$ defined by the equation $z_1 + \cdots + z_n = 2$. Let $\pi : \mathbb{R}^{n-1} \to \mathcal{H}$ denote the affine map defined by setting $\pi(z_1, \ldots, z_{n-1}) = (z_1, \ldots, z_{n-1}, 2 - (z_1 + \cdots + z_{n-1}))$. Then π is an affine isomorphism with $\pi(\mathbb{Z}^{n-1}) = \mathcal{H} \bigcap \mathbb{Z}^n$. Hence $\pi^{-1}(\mathcal{P}_G) \subset \mathbb{R}^{n-1}$ is an integral convex polytope with $\dim \pi^{-1}(\mathcal{P}_G) = n - 1$ and the toric ring $K[\pi^{-1}(\mathcal{P}_G)]$ is isomorphic to $K[\mathcal{P}_G]$ as homogeneous algebras over K.

Let δ denote the smallest integer for which the interior of $\delta(\pi^{-1}(\mathcal{P}_G))$ contains at least one integer point $(a_1, \ldots, a_{n-1})$. Since every connected component of $G_{[n]\setminus\{i\}}$ possesses at least one odd cycle, it follows from [8, Theorem 1.7 (a)] that the hyperplane defined by the equation $z_i = 0$ is a supporting hyperplane which defines a facet of $\mathcal{P}_G$ for each $1 \leq i \leq n$. Thus the hyperplane defined by the equation $z_i = 0$ is a supporting hyperplane which defines a facet of $\pi^{-1}(\mathcal{P}_G)$ for each $1 \leq i < n$. Thus by using [5, Corollary (1.2)] one has each $a_i = 1$ if $K[\pi^{-1}(\mathcal{P}_G)]$ is Gorenstein. If $\mathbf{e}_1 + \cdots + \mathbf{e}_{n-1}$ belongs to the interior of $\delta(\pi^{-1}(\mathcal{P}_G))$, then $\mathbf{e}_1 + \cdots + \mathbf{e}_{n-1} + q\mathbf{e}_n$ belongs to the interior of $\delta\mathcal{P}_G$ for some integer $q > 0$.

Since $K[\mathcal{P}_G]$ coincides with the Ehrhart ring of $\mathcal{P}_G$, it follows that there are edges $e_1, \ldots, e_m$ of G with $\mathbf{e}_1 + \cdots + \mathbf{e}_{n-1} + q\mathbf{e}_n = \rho(e_1) + \cdots \rho(e_m)$. Hence $q = 1$ and $\mathbf{e}_1 + \cdots + \mathbf{e}_n = \rho(e_1) + \cdots \rho(e_m)$. Thus G possesses a perfect matching with $\delta = n/2$ if $K[\pi^{-1}(\mathcal{P}_G)]$ is Gorenstein.

For a while, suppose that G possesses a perfect matching with $\delta = n/2$. Let $\mathcal{P}^\flat \subset \mathbb{R}^{n-1}$ denote the integral convex polytope $\delta(\pi^{-1}(\mathcal{P}_G)) - (\mathbf{e}_1 + \cdots + \mathbf{e}_{n-1})$. Then $\mathcal{P}^\flat$ is of standard type, i.e., $\dim \mathcal{P} = n - 1$ and the origin of $\mathbb{R}^{n-1}$ belongs to the interior of $\mathcal{P}$. Then [5, Corollary (1.2)] guarantees that the toric ring $K[\pi^{-1}(\mathcal{P}_G)]$ is Gorenstein if and only if the dual polytope [5, p. 631] of $\mathcal{P}^\flat$ is integral.

Now, by using [8, Theorem 1.7 (a)] again, it turns out that the equations of the supporting hyperplanes which defines the facets of $\mathcal{P}^\flat$ are the followings:

- $z_i = -1 \quad$ for each $\quad 1 \leq i < n$;
- $\sum_{i \in [n] \setminus (T \bigcup N(G;T))} z_i + 2\sum_{j \in N(G;T)} z_j = |T| - |N(G;T)| \quad$ if $\quad n \in T$;
- $2\sum_{i \in T} z_i + \sum_{j \in [n] \setminus (T \bigcup N(G;T))} z_j = |N(G;T)| - |T| \quad$ if $\quad n \in N(G;T)$;
- $\sum_{i \in T} z_i - \sum_{j \in N(G;T)} z_j = |N(G;T)| - |T| \quad$ if $\quad n \notin T \bigcup N(G;T)$,

where $T \neq \emptyset$ is a stable set of G for which the bipartite graph induced by T is connected and for which either $T \bigcup N(G;T) = [n]$ or every connected component of the induced subgraph $G_{[n] \setminus (T \bigcup N(G;T))}$ has at least one odd cycle.

Hence the dual polytope of $\mathcal{P}^\flat$ is integral if and only if (α) one has $|N(G;T)| = |T| + 1$ for each nonempty stable set T of G such that the bipartite graph induced by T is connected with $T \bigcup N(G;T) \neq [n]$ and that every connected component of $G_{[n] \setminus (T \bigcup N(G;T))}$ has at least one odd cycle and (β) one has $|T| = n/2 - 1$ for each stable set T of G such that the bipartite graph induced by T is connected with $T \bigcup N(G;T) = [n]$.

Consequently, when the toric ring $K[\mathcal{P}_G]$ is Gorenstein, the conditions (i), (ii) and (iii) are satisfied. Conversely, suppose that the conditions (i), (ii) and (iii) are satisfied. Since the hyperplane defined by the equation $z_i = 0$ is a supporting hyperplane which defines a facet of $\mathcal{P}_G$ for each $1 \leq i \leq n$, if $\gamma\mathcal{P}_G$, where $\gamma > 0$, contains at least one integer point $(a_1, \ldots, a_n)$, then each $a_i > 0$ and $\gamma \geq [(n+1)/2]$. It follows from (i), (ii) and (iii) that n is even and $\mathbf{e}_1 + \cdots + \mathbf{e}_n$ belongs to the interior of $(n/2)\mathcal{P}_G$. Thus the smallest number $\delta > 0$ for which $\delta\mathcal{P}_G$ contains at least one integer point is $\delta = n/2$ and $\mathbf{e}_1 + \cdots + \mathbf{e}_n$ belongs to the interior of $\delta\mathcal{P}_G$. Our discussion done already in the preceding paragraph guarantees that the toric ring $K[\mathcal{P}_G]$ is Gorenstein, as desired.

(a') In imitation of the preceding proof of (a) by using [8, Theorem 1.7 (b)] instead of [8, Theorem 1.7 (a)], one can easily give a proof of (a').

(b) The facets of the stable polytope $\mathcal{Q}_G$ is completely determined when G is a perfect graph ([4, Theorem 3.1]). In fact, when G is perfect, the equations of the supporting hyperplanes which defines the facets of $\mathcal{Q}_G$ are either $z_i = 0$ for $1 \leq i \leq n$ or $\sum_{W \subset [n]} z_i = 1$, where W is a maximal cliques of G. Let δ denote the smallest integer $\delta > 0$ for which the interior of $\mathcal{Q}_G$ contains at least one integer point. Then $\delta - 1$ coincides with the maximal cardinality of cliques of G and $\mathbf{e}_1 + \cdots + \mathbf{e}_n$ belongs to the interior of $\delta\mathcal{Q}_G$. It follows that the dual polytope of the integral polytope

$\delta\mathcal{Q}_G - (\mathbf{e}_1 + \cdots + \mathbf{e}_n) \subset \mathbb{R}^n$ of standard type is integral if and only if all maximal cliques of G have the cardinality $\delta - 1$. Hence [5, Corollary (1.2)] guarantees that the toric ring $K[\mathcal{Q}_G]$ is Gorenstein if and only if all maximal cliques have the same cardinality. $\square$

Example 2.2. The toric ring of the edge polytope of each of the finite connected graphs G_1 and G_2 drawn below is normal and Gorenstein.



G_1



G_2

3. Unimodal Gorenstein sequences

Let $\mathcal{P} \subset \mathbb{R}^N$ be a convex polytope. Recall that a $(q-1)$-simplex Σ each of whose vertices is a vertex of $\mathcal{P}$ is said to be a *special simplex* [2] in $\mathcal{P}$ if each facet of $\mathcal{P}$ contains exactly $q-1$ of the vertices of Σ.

Theorem 3.1. (a) *Let G be a finite connected graph as in Theorem 2.1 (a) or (a') and suppose that the toric ring $K[\mathcal{P}_G]$ of the edge polytope $\mathcal{P}_G$ of G is Gorenstein. Then there is a special simplex in $\mathcal{P}_G$.*

(b) *Let G be a perfect graph and suppose that the toric ring $K[\mathcal{Q}_G]$ of the stable polytope $\mathcal{Q}_G$ of G is Gorenstein. Then there is a special simplex in $\mathcal{Q}_G$.*

Proof. (a) Let $[n]$ be the vertex set of G. Since G possesses a perfect matching, it follows that $n = 2m$ is even and there exist m edges $e_1, \ldots, e_m$ of G with $\rho(e_1) + \ldots + \rho(e_m) = \mathbf{e}_1 + \cdots + \mathbf{e}_n$. Let Σ denote the $(m-1)$-simplex whose vertices are $\rho(e_1), \ldots, \rho(e_m)$. We claim that Σ is special in $\mathcal{P}_G$.

Theorem 2.1 together with [8, Theorem 1.7] give the complete information about the equations of the supporting hyperplanes which define the facets of the edge polytope $\mathcal{P}_G$. Let $\mathcal{H}_i$ denote the hyperplane defined by the equation $z_i = 0$. Then $\rho(e_j) \in \mathcal{H}_i$ if and only if $i \notin e_j$. Let $\mathcal{H}_T$ denote the hyperplane defined by the equation $\sum_{i \in T} z_i = \sum_{j \in N(G;T)} z_j$, where T is a stable set of G, which is the supporting hyperplane of a facet of $\mathcal{P}_G$. Let C denote the set of those $1 \leq i \leq m$ with $\rho(e_i) \bigcap T \neq \emptyset$ and D the set of those $1 \leq i \leq m$ with $\rho(e_i) \bigcap N(G;T) \neq \emptyset$. In either the case of $|N(G;T)| = |T| + 1$ or the case of $|N(G;T)| - 1 = |T| + 1 = m$, one has $C \subset D$ with $|C| = |D| - 1$. Let $i_0 \in D \setminus C$. Then $\rho(e_j) \in \mathcal{H}_T$ if and only if $j \neq i_0$. Thus Σ is special in $\mathcal{P}_G$ as desired.

(b) Let G be a perfect graph on $[n]$ and suppose that all maximal cliques have the cardinality q. Since G is perfect, the chromatic number of G is equal to q. Thus there is a map $\varphi : [n] \to [q]$ with the property that $\varphi(i) \neq \varphi(j)$ if $\{i,j\} \in E(G)$. Let W'_ℓ denote the stable set $\{i \in [n] : \varphi(i) = \ell\}$ for each $1 \leq \ell \leq q$. We assume that $\mathcal{Q}_G$ is not a simplex. Thus one of the stable sets $W'_1, \ldots, W'_q$ contains at least two vertices. Let, say, W'_1 contain at least two vertices and fix $i_0 \in W'_1$. Let $W_0 = \{i_0\}, W_1 = W'_1 \setminus \{i_0\}$ and $W_\ell = W'_\ell$ for $2 \leq \ell \leq q$. Each of $W_0, W_1, \ldots, W_q$ is a stable set of G and $\sum_{\ell=0}^{q} \rho(W_\ell) = \mathbf{e}_1 + \cdots + \mathbf{e}_n$. Let Σ denote the q-simplex with $q+1$ vertices $\rho(W_0), \rho(W_1), \ldots, \rho(W_q)$. We claim that Σ is special in $\mathcal{Q}_G$.

Recall that the equation of the supporting hyperplanes which defines the facets of the stable polytope of G are either (i) $x_i = 0$ for $1 \leq i \leq n$ or (ii) $\sum_{i \in W} x_i = 1$, where W is a maximal clique of G. If F_i is the facet defined by $x_i = 0$, then $\rho(W_\ell) \in F_i$ if and only if $i \notin W_\ell$. Since $[n]$ is the disjoint union $W_0 \bigcup W_1 \bigcup \cdots \bigcup W_q$, it follows that F_i contains exactly q of the vertices of Σ. Let F'_W denote the facet defined by $\sum_{i \in W} x_i = 1$, where W is a maximal clique of G. Since each of the subsets $W \cap (W_0 \bigcup W_1), W \cap W_2, \ldots, W \cap W_q$ of $[n]$ consists of one element, it follows that each of the vertices $\rho(W_2), \ldots, \rho(W_q)$ belongs to F'_W and that $\rho(W_0) \in F'_W$ (resp. $\rho(W_1) \in F'_W$) if and only if $i_0 \in W$ (resp. $i_0 \notin W$). Hence F'_W contains exactly q of the vertices of Σ. □

By using Example 1.1 together with [16, Theorem 3.2], it turns out that the above Theorem 3.1 (b) generalize Reiner–Welker [12, Corollary 3.8].

Now, by virtue of [2, Theorem 3.5], one has a rich class of unimodal Gorenstein sequences [17, p. 66]. It is known [11, Example 1.3 (c)] the stable polytope of a perfect graph is compressed.

The edge polytope of a finite connected graph is unimodular, i.e., all of its triangulations are unimodular, if and only if any two odd cycles of G possess at least one common vertex. In particular the edge polytope of a finite connected bipartite graph is unimodular. The edge polytope of G_1 of Example 2.2 is compressed ([10]) but not unimodular, and that of G_2 is unimodular. A combinatorial characterization of finite graphs G for which the edge polytope $\mathcal{P}_G$ is compressed is given in [7, Theorem 4.1].

Corollary 3.2. (a) *Let G be a finite connected graph as in Theorem 2.1 (a) and suppose that the edge polytope $\mathcal{P}_G$ is compressed and that the toric ring $K[\mathcal{P}_G]$ is Gorenstein. Then the h-vector of $K[\mathcal{P}_G]$ is unimodal.*

(a') *Let G be a finite 2-connected bipartite graph and suppose that the toric ring $K[\mathcal{P}_G]$ is Gorenstein. Then the h-vector of $K[\mathcal{P}_G]$ is unimodal.*

(b) *Let G be a perfect graph and suppose that the toric ring $K[\mathcal{Q}_G]$ of the stable polytope $\mathcal{Q}_G$ of G is Gorenstein. Then the h-vector of $K[\mathcal{Q}_G]$ is unimodal.*

We conclude the present paper with

Example 3.3. Let $n \geq 3$ and G the finite connected graph on $[2n]$ drawn below. Let n be odd. (If n is even, then $K[\mathcal{P}_G]$ is not Gorenstein by Theorem 2.1 (a').) By virtue of [7, Theorem 4.1] it turns out that the edge polytope $\mathcal{P}_G$ of G is compressed. By using Theorem 2.1 (a) it follows that the toric ring $K[\mathcal{P}_G]$ is (normal and) Gorenstein. Moreover, we can compute the h-vector explicitly. Since the graph G satisfies the condition in [9, Theorem 1.2], the "toric ideal" I_G of G is generated by quadratic binomials which correspond to even cycles of G of length 4. There exists a reverse lexicographic order such that the initial monomials of the quadratic binomials are relatively prime. Since the set of quadratic binomials is a Gröbner basis of I_G with respect to $<_{rev}$, it follows that the initial ideal is generated by n monomials which are squarefree, quadratic and relatively prime. Thus the h-vector of $K[\mathcal{P}_G]$ is $\left(1, n, \binom{n}{2}, \cdots, \binom{n}{n-2}, n, 1\right)$.



References

[1] C. Berge, Some classes of perfect graphs, *in* "Graph Theory and Theoretical Physics" (F. Harary, Ed.), Academic Press, London, 1967, pp. 155 – 165.

[2] C. A. Athanasiadis, Ehrhart polynomials, simplicial polytopes, magic squares and a conjecture of Stanley, preprint, 2003.

[3] W. Bruns and J. Herzog, "Cohen–Macaulay Rings," Cambridge University Press, Cambridge, New York, Sydney, 1993.

[4] V. Chvátal, On certain polytopes associated with graphs, *J. Combin. Theory* (B) **18** (1975), 138 – 154.

[5] E. De Negri and T. Hibi, Gorenstein algebras of Veronese type, *J. Algebra* **193** (1997), 629 – 639.

[6] T. Hibi, "Algebraic Combinatorics on Convex Polytopes," Carslaw Publications, Glebe, N.S.W., Australia, 1992.

[7] H. Ohsugi, Unimodular regular triangulations of $(0, 1)$-polytopes associated with finite graphs, *in* "Algebraic Engineering" (C. L. Nehaniv and M. Ito, Eds.), World Scientific, Singapore, 1999, pp. 159 – 171.

[8] H. Ohsugi and T. Hibi, Normal polytopes arising from finite graphs, *J. Algebra* **207** (1998), 409 – 426.

[9] H. Ohsugi and T. Hibi, Toric ideals generated by quadratic binomials, *J. Algebra* **218** (1999), 509 – 527.

[10] H. Ohsugi and T. Hibi, Compressed polytopes, initial ideals and complete multipartite graphs, *Illinois J. Math.* **44** (2000), 391 – 406.

[11] H. Ohsugi and T. Hibi, Convex polytopes all of whose reverse lexicographic initial ideals are squarefree, *Proc. Amer. Math. Soc.* **129** (2001), 2541 – 2546.

[12] V. Reiner and V. Welker, On the Charney–Davis and Neggers–Stanley conjectures, preprint, 2002.

[13] A. Simis, W. V. Vasconcelos, and R. H. Villarreal, The integral closure of subrings associated to graphs, *J. Algebra* **199** (1998), 281 – 289.

[14] R. P. Stanley, The number of faces of a simplicial convex polytope, *Adv. in Math.* **35** (1980), 236 – 238.

[15] R. P. Stanley, Decompositions of rational convex polytopes, *Ann. Discrete Math.* **6** (1980), 333 – 342.

[16] R. P. Stanley, Two poset polytopes, *Disc. Compt. Geom.* **1** (1986), 9 – 23.

[17] R. P. Stanley, "Combinatorics and Commutative Algebra," Second Ed., Birkhäuser, Boston, 1996.

[18] B. Sturmfels, "Gröbner Bases and Convex Polytopes," Amer. Math. Soc., Providence, RI, 1995.

Hidefumi Ohsugi
Department of Mathematics
Faculty of Science
Rikkyo University
Toshima, Tokyo 171–8501, Japan
E-mail:ohsugi@rkmath.rikkyo.ac.jp

Takayuki Hibi
Department of Pure and Applied Mathematics
Graduate School of Information Science and Technology
Osaka University
Toyonaka, Osaka 560–0043, Japan
E-mail:hibi@math.sci.osaka-u.ac.jp